

Women, Sex, Gender and Dementia Cross-Cutting Program: Phase II Publications

D'Amico D, Parrott MD, Greenwood CE, Ferland G, Gaudreau P, Belleville S, Laurin D, Anderson ND, Kergoat M, Morais JA, Presse N, Fiocco AJ (2020). **Sex differences in the relationship between dietary pattern adherence and cognitive function among older adults: Findings from the NuAge study.** *BMC Nutrition Journal*, 19(1):58.
<https://nutritionj.biomedcentral.com/articles/10.1186/s12937-020-00575-3>

Gervais NJ, Au A, Almey A, Duchesne A, Gravelins L, Brown A, Reuben R, Baker-Sullivan E, Schwartz DH, Evans K, Bernardini MQ, Eisen A, Meschino WS, Foulkes WD, Hampson E, Einstein G (2020). **Cognitive markers of dementia risk in middle-aged women with bilateral salpingo-oophorectomy prior to menopause.** *Neurobiology of Aging*, 94, 1-6.
<https://doi.org/10.1016/j.neurobiolaging.2020.04.019>.

McFall, GP, Bäckman L, Dixon RA (2019) .**Nuances in Alzheimer's Genetic Risk Reveal Differential Predictions of Non-demented Memory Aging Trajectories: Selective Patterns by APOE Genotype and Sex.** *Current Alzheimer Research*. 16(4):302-315.
<https://pubmed.ncbi.nlm.nih.gov/30873923/>

Sourial N, Vedel I, Godard-Sebillotte C, Etches J, Arsenault-Lapierre G, Bronskill SE (2020). **Sex Differences in Dementia Primary Care Performance and Health Service Use: A Population-Based Study.** *Journal American Geriatrics Society*. 68(5):1056-1063.
<https://onlinelibrary.wiley.com/doi/abs/10.1111/jgs.16347>

Thibeau S, McFall GP, Camicioli R, Dixon, RA (2019). **Physical Activity and Mobility Differentially Predict Nondemented Executive Function Trajectories: Do Sex and APOE Moderate These Associations?.** *Gerontology*. 65(6):640-648.
<https://www.karger.com/Article/Pdf/496442>

Trenaman SC, Rideout M, Andrew MK (2019). **Sex and gender differences in polypharmacy in persons with dementia.** *SAGE Open Medicine*. 22;7:2050312119845715.
<https://pubmed.ncbi.nlm.nih.gov/31041100/>

Ursenbach J, O'Connell ME, Kirk A, Morgan D (2020). **Evidence for measurement bias of the short form health survey based on sex and metropolitan influence zone in a secondary care population.** *Health and Quality of Life Outcomes*. 18(1):91.
<https://hqlo.biomedcentral.com/articles/10.1186/s12955-020-01318-y>